

# **Morphodynamics and Geology of the Southeastern Virginia Shelf: False Cape Shoals Area (Phase 2)**

Stephen P. Murray, Ph.D.  
Physical Oceanography  
Code 322PO

Randolph A. McBride, Ph.D.  
Assistant Professor of Geology  
Environmental Science and Policy, MS 5F2  
George Mason University  
Fairfax, VA 22030

Phone: 703-993-1210 Fax: 703-993-1216 E-mail: [rmcbride@gmu.edu](mailto:rmcbride@gmu.edu)

Award Number: N00014-00-1-0247

## **LONG-TERM GOALS**

The long-term goal is to investigate the morphodynamics of shore-oblique shoals at False Cape, Virginia from geologic and hydrodynamic perspectives to address sand body genesis, geometry, and evolution. An additional goal of this ROPO is to continue analysis and publication of data taken in Bab el Mandab, the entrance Strait to the Red Sea.

## **OBJECTIVES**

Geologic data (i.e., sediment cores) will be collected to analyze the geologic framework (i.e., sedimentology, stratigraphy) of the shore-oblique shoals at False Cape, Virginia.

## **APPROACH**

Shore-oblique shoals along the southeastern Virginia shelf have proven to be viable sources of sand for beach replenishment. Numerous studies have addressed the distribution and orientation of shore-oblique sand shoals along the U.S. Atlantic shelf (e.g., Duane et al., 1972; Swift et al., 1972; Swift and Field, 1980; Figueiredo et al., 1981; McBride and Moslow, 1991) but more detailed geologic data are needed to characterize the internal stratigraphy and sedimentology (e.g., physical and biological sedimentary structures, grain size analysis, foraminiferal analysis) in order to document shoal genesis, evolution, and migration (see Snedden and Dalrymple, 1999; Snedden et al., 1999; Dyer and Huntley, 1999). Huthnance (1982a, b) has proposed a purely fluid dynamical model to explain the flow field responsible for the morphodynamics of shore-oblique shoals but this model needs further testing using actual field data (geologic and hydrodynamic information). This ROPO project helps to address these issues.

## **WORK COMPLETED**

In June 2001, a total of 20 vibracores (sediment cores) were collected in the False Cape, VA/NC area. Fourteen of the cores were collected from the continental shelf directly on or adjacent to False Cape

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2001</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2001 to 00-00-2001</b>	
4. TITLE AND SUBTITLE <b>Morphodynamics and Geology of the Southeastern Virginia Shelf: False Cape Shoals Area (Phase 2)</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Environmental Science and Policy, MS 5F2,,George Mason University,,Fairfax,,VA, 22030</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>The long-term goal is to investigate the morphodynamics of shore-oblique shoals at False Cape, Virginia from geologic and hydrodynamic perspectives to address sand body genesis, geometry, and evolution. An additional goal of this ROPO is to continue analysis and publication of data taken in Bab el Mandab, the entrance Strait to the Red Sea.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

shoal, whereas six cores were drilled on the adjacent barrier island at Old Currituck inlet (i.e., flood-tidal delta). Athena Technologies, Inc. from Columbia, SC was subcontracted to provide coring vessels, equipment, and support.

In the open ocean, the RV *Artemis* was maneuvered to the desired vibracore sample site using a Trimble Ag132 DGPS. Once on station the vibracoring system was prepared for sampling. The sample barrel was lowered to the seafloor by attaching lengths of drill stem until the seafloor was reached. Once the end of the core barrel touched bottom, the vibracore was turned on. When penetration reached refusal, the vibracore was turned off. The sample barrel was then winched to the deck of the RV *Artemis*, where the vibracores were cut, capped, and measured for recovery. Fourteen vibracores, ranging from 2 to 6 m, were collected from the continental shelf using the RV *Artemis*.

Six vibracore samples were taken in Currituck Sound on an abandoned flood-tidal delta. These cores ranged in length from 2 to 8 m and were taken using Athena's terrestrial sampling gear. This gear included a backpack mounted vibracore system, an aluminum tripod, and a hand winch. This vibracore process used 3-inch diameter aluminum sample barrels instead of the 2 3/8 inch galvanized. Athena used a Jon Boat to carry the equipment to the desired sample locations that were located in the field using a DGPS provided by George Mason University. Once on station the equipment was carried a short distance to the desired site. The vibrator head was attached to the sample barrel and raised to the vertical position. The vibracore machine was then turned on and allowed to vibrate in until refusal. The sample barrel was then capped to provide a seal at the top, the tripod positioned over the sample barrel, and the hand winch was then attached to the tripod. The sample was then winched out of the ground, cut down to the proper size, and capped on both ends. Each vibracore sample was sectioned into one-meter lengths and each was cut open longitudinally. These were then placed into plastic tubing to preserve the sample and transported back to George Mason University for analysis.

Dr. Randy McBride is now processing and describing the sediment cores in the Coastal Geology Laboratory at George Mason University. In addition, a Ph.D. student in Environmental Science and Policy, who is working under the direction of Dr. McBride, is performing a micropaleontological study of the foraminiferal assemblages in the cores for her dissertation topic. Identifying the foraminiferal assemblages will play a major role in understanding the stratigraphy and geologic processes characterizing the shoal sand bodies.

## **RESULTS**

Oceanographic coring must be done in the summer months when sea conditions are optimal for drilling operations. As a result, data collection had to be postponed eight months to June 2001 in order to ensure appropriate sea conditions. As such, data analysis on the sediment cores has only recently begun and no results are available at this time.

## **RELATED PROJECTS**

This ROPO project complements a recently awarded research project to George Mason University (GMU) entitled "Hard Mineral Resources of the Southeastern Virginia Shelf: Preliminary Geologic Investigation of the False Cape Shoals Area" (Principal Investigator: R. A. McBride). This latter GMU project was funded by an ongoing cooperative agreement between the Minerals Management Service (MMS), US Dept. of the Interior and the Virginia Institute of Marine Science (VIMS) of the College of William and Mary. This federal/state cooperative agreement is for the purpose of

conducting geological and physical oceanographic studies relating to the identification, characterization, and potential use of offshore sand resources for beach nourishment along coastal Virginia.

## REFERENCES

- Duane, D.B., Field, M.E., Meisburger, E.P. Swift, D.J.P., and Williams, S.J., 1972. Linear shoals on the Atlantic inner continental shelf, Florida to Long Island. In: Swift et al. (eds.), *Shelf Sediment Transport: Process and Pattern*. Dowden, Hutchinson, and Ross, Stroudsburg, PA, pp. 447-498.
- Dyer, K.R. and Huntley, D.A., The origin, classification, and modeling of sand banks and ridges. *Continental Shelf Research*, 19:1285-1330.
- Figueiredo, A.G., Swift, D.J.P., Stubblefield, W.L., and Clarke, T.L., 1981. Sand ridges on the inner Atlantic shelf of North America: morphometric comparisons with Huthnance stability model. *Geo-Marine Letters* 1:187-191.
- Huthnance, J.M., 1982a. On one mechanism forming linear sand banks. *Estuarine, Coastal Shelf Science*, 14:79-99
- Huthnance, J.M., 1982b. On the formation of sandbanks of finite extent. *Estuarine, Coastal Shelf Science*, 15:277-299.
- McBride, R. A. and T. F. Moslow, 1991. Origin, evolution, and distribution of shoreface sand ridges, Atlantic inner shelf. USA. *Marine Geology*, 97:57-85.
- Snedden, J.W. and Dalrymple, R.W., 1999, Modern shelf sand ridges: from historical perspective to a unified hydrodynamic and evolutionary model, In: Bergman, K.M. and Snedden, J.W. (editors), *Isolated Shallow Marine Sandbodies: Sequence Stratigraphic Analysis and Sedimentologic Interpretation*: Tulsa, OK, Society of Sedimentary Geology (SEPM) Special Publication #64, pp. 13-28.
- Snedden, J.W., Kreisa, R.D., Tillman, R.W., Culver, S.J, and Schweller, W.J., 1999, An expanded model for modern shelf sand ridge genesis and evolution on the New Jersey Atlantic shelf, In: Bergman, K.M. and Snedden, J.W. (editors), *Isolated Shallow Marine Sandbodies: Sequence Stratigraphic Analysis and Sedimentologic Interpretation*: Tulsa, OK, Society of Sedimentary Geology (SEPM) Special Publication #64, pp. 147-163.
- Swift, D.J.P. and Field, M.E., 1981. Evolution of a classic sand ridge field: Maryland sector, North American inner shelf. *Sedimentology*, 28:461-482.
- Swift, D.J.P., Kofoed, J.W., Saulsbury, F.P., and Sears, P., 1972. Holocene evolution of the shelf surface, central and southern Atlantic shelf of North America. In: Swift et al. (eds.), *Shelf Sediment Transport: Process and Pattern*. Dowden, Hutchinson, and Ross, Stroudsburg, PA, pp. 447-498.